

MORNINGSIDE DENTAL CARE

After Sedation instructions for Companion & Patient

- 1) PATIENT CANNOT DRIVE FOR 24 HOURS AFTER TAKING SEDATION MEDICATION
- 2) DO NOT OPERATE ANY HAZARDOUS DEVICES FOR 24HOURS
- 3) A RESPONSIBLE PERSON SHOULD BE WITH THE PATIENT UNTIL HA/SHE HAS FULLY RECOVERED FROM THE EFFECTS OF THE SEDATION.
- 4) PATIENT SHOULD NOT GO UP AND DOWN STAIRS UNATTENDED. LET THE PATIENT STAY ON THE GROUND FLOOR UNTIL RECOVERED.
- 5) HAVING NUTRITION AFTER SEDATION IS IMPORTANT. THE PATIENT SHOULD BEGIN EATING APPROPRIATE FOODS AS SOON AS POSSIBLE. DO NOT DELAY.
- 6) PATIENT NEEDS TO DRINK PLENTY OF FLUIDS AS SOON AS POSSIBLE.
- 7) PATIENT MAY SEEM ALERT WHEN HE/SHE LEAVES. THIS MAY BE MISLEADING SO DO NOT LEAVE THE PATIENT ALONE.
- 8) ALWAYS HOLD PATIENTS ARM WHEN WALKING.
- 9) CALL US IF YOU HAVE ANY QUESTIONS OR DIFFICULTIES. IF YOU FEEL THAT YOUR SYMPTOMS WARRANT A PHYSICIAN AND YOU ARE UNABLE TO REACH US, GO TO THE CLOSEST EMERGENCY ROOM IMMEDIATELY.
- 10) PATIENT SHOULD NOT CARRY, SLEEP NEXT TO OR BE LEFT ALONE WITH YOUNG CHILDREN FOR A PERIOD OF NO LESS THAN 24 HOURS AFTER THE LAST DOSAGE OF MEDICATION.
- 11) DRIVE DIRECTLY HOME. CALL THE OFFICE WHEN YOU ARRIVE AT HOME AND THE PATIENT IS COMFORTABLE AND SAFE.

FOLLOWING MOST DURGICAL PROCEDURES THERE MAY OR MAY NOT BE PAIN, DEPENDIDN ON YOUR THRESHHOLD FOR PAIN. YOU WILL BE PROVIDED WITH MEDICATION FOR DISCOMFORT THAT IS APPROPRIATE FOR YOU. IN MOST CASES A NON-NARCOTIC PAIN REGIMEN WILL BE GIVEN CONSISTING OF *ACETAMINOPHEN* (TYLENOL) AND *IBUPROFEN* (ADVIL). THESE TWO MEDICATIONS **TAKEN TOGETHER**, WILL BE AS EFFECTIVE AS A NARCOTIC WITHOUT ANY OF THE SIDE EFFECTS ASSOCIATED WITH NARCOTICS. IF A NARCOTIC HAS BEEN PRESCRIBED, FOLLOW THE DIRECTIONS CAREFULLY. IF YOU HAVE ANY QUESTIONS ABOUT THESE MEDICATIONS INTERACTING WITH OTHER MEDICATIONS YOU ARE PRESENTLY TAKING, PLEASE CALL OUR OFFICE, YOUR PHYSICIAN AND/OR YOUR PHARMACIST.

DR. WILLIAMS HOME: 646-961-3513

DATE:

COMPANIONS SIGNATURE:

PATIENTS SIGNATURE:

MEDICATIONS: Take only when checked
___ AMOXICILLIN- Fill prescription and take s directed
___ ERYTHROMYCIN- Fill prescription and take as directed
___ TYLENOL (ACETAMINOPHEN)- take two every six hours
___ ADVIL (IBUPROFEN) - Take two every six hours
___ LORATAB- For PAIN ONLY Take one every 6 hours
___ VITAMIN C- one (1000mg) at every meal 3x a day.
___ Co Q 10- 50mg 2X a day